

How to Test Substances for Fentanyl

You will need:







1 tsp (5mL) of water which is 1 water bottle cap

Substance you would like to test

Fentanyl Test Strip



How to Test Substances for Fentanyl

You will need:







1 tsp (5mL) of water which is 1 water bottle cap

Substance you would like to test

Fentanyl Test Strip



Adjusting to test stimulants:

When testing stimulants for fentanyl, use a 1/4 cup of water (which is equivalent to 1 full shot glass or approximately 1/8 of a standard water bottle) to prevent false positives from the test strip.




Adjusting to test stimulants:


When testing stimulants for fentanyl, use a 1/4 cup of water (which is equivalent to 1 full shot glass or approximately 1/8 of a standard water bottle) to prevent false positives from the test strip.

Instructions:

- Crush a small amount of substance (skip if already crushed).
- Measure out 5-10mg of the crushed substance: enough to cover Lincoln's hair on a penny.
 - No penny? No problem! Measure substance by covering the black circle with your substance and test that amount.
- Pour your water into your clean container or cup. Put the measured substance in the water and swirl it until it is completely dissolved.
- Hold the solid blue end of the test strip and dip the wavy end into the water. Don't dip past the last line. Hold in water for 10 seconds.

How to Read the Strip






Instructions:

- Crush a small amount of substance (skip if already crushed).
- Measure out 5-10mg of the crushed substance: enough to cover Lincoln's hair on a penny.
 No penny? No problem! Measure substance by covering the black circle with your substance and test that amount.
- Pour your water into your clean container or cup. Put the measured substance in the water and swirl it until it is completely dissolved.
- Hold the solid blue end of the test strip and dip the wavy end into the water. Don't dip past the last line. Hold in water for 10 seconds.

How to Read the Strip

Take the strip out of the water and wait 1 minute for your results



Use a bright light to look at the strips and count the number of red lines. Even, light red lines count!

This project has been funded by grant number NH28CE003534 from the Centers for Disease Control. The contents are those of the author(s) and do not necessarily represent the official views of, nor an endorsement, by the CDC, or the U.S. Government.

This project has been funded by grant number NH28CE003534 from the Centers for Disease Control. The contents are those of the author(s) and do not necessarily represent the official views of, nor an endorsement, by the CDC, or the U.S. Government.

Harm Reduction Tips

that you engage in additional harm reduction practices, while using: such as those listed below, to reduce the risk of harm accurate. When using substances it is recommended No drug is completely safe and no test is 100%



people you trust or call the Never Never use alone! Always use with before using the substance Test for xylazine and fentanyl

Always carry naloxone on you and administer it in the event of an (877) 696-1996 **Use Alone Hotline at**

overdose

test strips, contact: For help or to get more

SNHD Linkage to Action Phone: 702-759-1381 and press 2 Email: L2A@snhd.org

Harm Reduction Tips

such as those listed below, to reduce the risk of harm that you engage in additional harm reduction practices, accurate. When using substances it is recommended No drug is completely safe and no test is 100%



- Test for xylazine and fentanyl Never use alone! Always use with before using the substance
- (877) 696-1996 people you trust or call the Never **Use Alone Hotline at**
- overdose administer it in the event of an Always carry naloxone on you and

test strips, contact: For help or to get more

SNHD Linkage to Action Phone: 702-759-1381 and press 2 Email: L2A@snhd.org

Matter You

Stop by and get more

supplies

Matter You

supplies Stop by and get more